VOLUME 04 ISSUE 01 Pages: 36-40

SJIF IMPACT FACTOR (2021: 5.478) (2022: 5.636) (2023: 6.741)

OCLC - 1368736135















Copyright: Original content from this work may be used under the terms of the creative commons attributes 4.0 licence.



Research Article

STUDY OF ENTOMOPATHOGENIC NEMATODES HISTORY

Submission Date: January 01, 2024, Accepted Date: January 03, 2024,

Published Date: January 05, 2024

Crossref doi: https://doi.org/10.37547/ijasr-04-01-06

Kurbonova Nafosat Sattar Kizi

Research Institute Of Plant Quarantine And Protection, Uzbekistan

Tillyakhodjayeva Nigora Ruzimatovna

Research Institute Of Plant Quarantine And Protection, Uzbekistan

Karimova Rikhsiniso Miratxamovna

Research Institute Of Plant Quarantine And Protection, Uzbekistan

ABSTRACT

Entomopathogen nematodes Uzbekistan fauna for yet not studied species this is for the first time in the article of Uzbekistan potatoes from agrobiocenoses separate received entomopathogenic of nematodes morpho-biological features the world scientists in experience to be studied ie entomopathogenic of nematodes study history about analytical data present will be done.

Keywords

Entomopathogen nematodes, Uzbekistan, potatoes, Steinernema, Heterorhabditis, infectious juvenil stage.

INTRODUCTION

Entomopathogenic nematodes feed on the internal tissues of the insect with the help of bacteria that live symbiotically in the intestine, develop and reproduce. The term "Entomopathogen" also means "entomos" - insect, "pathogen" - disease-causing,

disease-causing insect. Representatives of the Steinernematidae and Heterorhabditidae families entomopathogenic nematodes pathogenicity in more than 200 types of pest insects belonging to 100 genera in symbiosis with bacteria of Xenorhabdus and Photorhabdus

VOLUME 04 ISSUE 01 Pages: 36-40

SJIF IMPACT FACTOR (2021: 5.478) (2022: 5.636) (2023: 6.741)

OCLC - 1368736135











genera, respectively, and kill their host within 2 days.

The first entomopathogenic nematodes were described by Steiner in 1923 as Aplectana kraussei (now Steinernema kraussei), but this information was based on very little scientific evidence.

Neoaplectana glaseri, was later reported by Glaser and Fox in 1930 . Steiner includes this species in the Oxyuridae family.

Only Yaroslav Weiser isolated and described Neoaplectana carpocapsae moth larvae in Europe. Also, in 1955, Dutki and Hough isolated the steinernematid strain DD-136 from moth larvae in the eastern part of North America, after which serious research on the pathogenicity and development of entomopathogenic nematodes began.

In 1965, with the help of morphological and physiological studies, Poinar noted that Weiser's S. carpocapsae Czechoslovak strain and the North American DD-136 nematode are specific.

in 1965 Symbiosis with S. carpocapsae by Poinar and Thomas associated symbiotic bacterium is described.

Later, in 1966 - 1977, Poinar and Leutenegger described the location of bacteria in nematode larvae in the infectious stage using light and electron microscopy.

Poinar and Thomas, 1967, identified the role of bacteria in nematode developmental stages and killing insect pests and reported a new bacterial family, Xenorhabdus.

The genus Heterorhabditis was first described by Poinar in 1976, and its symbiotic bacterium Xenorhabditis luminescens was described by Thomas and Poinar in 1979. The luminescence ability of this symbiotic bacterium was so strong that all infected insect corpses glowed in the dark.

Later, in 1903, Boemari declared this bacterial species to be a representative of the genus Photorhabdus. The mechanism of action of bacteria was clarified by Milstead in 1977.

According to genetic analyzes conducted by Kiontke in 2007, the genus Heterorhabditis is considered a close group to vertebrate parasitic nematodes, and both groups diverged independently from the free-living Rhabditis group.

The mutualistic existence of representatives of the genus Heterorhabditis with a unique group of luminescent symbiotic bacteria, its ability to enter the body of healthy insects, the alternation of sexual and hermaphrodite generations, and its unique morphology explain the fact that these species are given the status of a family.

Another feature not observed in Steinernema and other rhabditids is the presence of a dorsal "hook" on the tip of the head of third instar larvae of Heterorhabditis. Bedding and Molyneux (1982) noted in their research that this structure allows the potential host to enter its body cavity through the outer tegument or trachea and intestinal walls.

37

VOLUME 04 ISSUE 01 Pages: 36-40

SJIF IMPACT FACTOR (2021: 5.478) (2022: 5.636) (2023: 6.741)

OCLC - 1368736135











According to Akhurst and Boemare's research, all Steinernema species have a symbiotic relationship with Xenorhabdus bacteria, and Heterorhabditis species have a symbiotic relationship with Photorhabdus bacteria. It was the discovery of the symbiotic relationship between entomopathogenic nematodes and bacteria that marked a turning point in the use of nematodes as commercial biological control agents.

In 1959, Dutky was the first to realize the antibiotic properties of bacteria living in symbiosis with S. carpocapsae, when he determined the role of symbiotic bacteria in the reproduction of entomopathogenic nematodes developing inside the insect corpse.

Several antibiotics, including xenorhabdins, xenocaucins, hydroxystilbenes, indole anthraquinone derivatives, were subsequently Xenorhabdus synthesized from Photorhabdus cultures in 2002 by Webster et al.

Primary stage larvae of entomopathogenic nematodes grow maximally and carry out antibiotic production. But the primary phase goes directly to the second phase, in which nematode growth slows down and antibody production stops. Poinar noted in 1980 that this process is a major obstacle to the commercial use of nematodes.

In turn, insects have many protective reactions against entomopathogenic nematodes, and the most important of these are melanization and encapsulation. Usually, the bacteria are able to kill the insect much faster than the insect's defense

mechanism work. can However. some experimental host organisms, such as flies, perform a very rapid melanization reaction that kills the nematodes before they can activate the symbiotic bacteria. Through this research, Bronskill and Welch presented many new scientific conclusions in 1962.

Furthermore, the fact that the infective larvae of S. carpocapsae can be easily destroyed even by Galleria mellonella. the most common experimental organism used for the propagation of entomopathogenic nematodes, if they do not have symbiotic bacteria, was demonstrated in Poinar's 1969 research work, announced.

Also, Veremchuk and Issi, 1970, natural combinations of nematodes in the infectious stage, for example, unicellular and fungal representatives, also quickly destroy nematodes in the insect organism. For example, when moths primarily infected with microsporidia are infected by S. carpocapsae, they note that the microsporidia also infect nematodes.

Poinar found that naturally occurring populations of entomopathogenic nematodes can also be infected by microsporidia, whose infective larval stages are susceptible to infection by several different common soil fungi. emphasizes the need to test for suspected nematophagous fungi.

REFERENCES

1. Campos-Herrera, R., et al., Unraveling the intraguild competition between Oscheius spp. nematodes and entomopathogenic

Volume 04 Issue 01-2024

38

VOLUME 04 ISSUE 01 Pages: 36-40

SJIF IMPACT FACTOR (2021: 5.478) (2022: 5.636) (2023: 6.741)

OCLC - 1368736135











- nematodes: Implications for their natural distribution in Swiss agricultural soils. I Inverteb Pathol, 2015. 132: p. 216-227.
- 2. Campos-Herrera,R.,et al., Intraspecific virulence entomopathogenic of nematodes against the pests. J Nematol, 2021.53.
- 3. Dillman , AR, et al., jumps with greater velocity and acceleration than previously reported. Micropubl Biol, 2021. 2021.
- Dillman, AR, et al., Comparative genomics 4. of Steinernema reveals deeply conserved gene regulatory networks. Genome Biol, 2015. 16: p. 200.
- Dutka, A., A. McNulty, and SM Williamson, 5. A new threat to bees? Entomopathogenic nematodes used in biological pest control cause rapid mortality in Bombus terrestris . Peerl , 2015. 3 : p. e1413.
- Filipjev , IN (1934). 6. Miscellanea Nematologica 1. Eine neue Art der Gattung Neoaplectana Steiner nebst Bermerkungen über die systematische Stellung der letzteren . Magazine de Parasitologie du Musée Zoologique de l'Académie des Sciences de l'USSR 4, 229-240.
- 7. GLASER, RW (1932).Studies on Neoaplectana glaseri, a nematode parasite of the Japanese beetle (Popillia japonica). New Jersey Department of Agriculture Circular 211.
- 8. Glaser, RW. & Farrell, CC (1935). Field experiments with the Japanese beetle and its nematode parasite. Journal of the New York Entomological Society 43, 345-371.

- 9. Hominick, WM, Briscoe, BR, Del Pino, FG, Heng, IA, Hunt, DJ, Kozodoy, E., Mráïc Ek, Z., Nguyen, KB, Reid, AP, Spiridonov, SE, Stock, P., Sturhan , D., Waturu , C. & Yoshida, M. (1997). Biosystematics of entomopathogenic nematodes: current status, protocols and de. nitions. Journal of Helminthology 71, 271-298.
- 10. Kaya, HK & Stock, SP (1997). Techniques in insect nematology. In: Lacey, LA (Ed.) Techniques in insects pathology. London, Academic Press, pp. 281-324.
- 11. Kaplan, F., et al., Conspecific heterospecific pheromones stimulate dispersal of entomopathogenic nematodes during quiescence. Sci Rep, 2020. 10 (1): p. 5738.
- 12. Lacey, LA & Chauvin, RL (1999). Entomopathogenic nematodes for control of diapausing codling moth (Lepidoptera: Tortricidae) in fruit bins. Journal of Economic Entomology 92, 104-109
- Nadler, SA, Adams, BJ, Lyons, ET, Delong, **13**. RL & Melin, SR (2000). Molecular and morphometric evidence for separate species of Uncinaria (Nematoda : Ancylostomatidae) in California sea lions and Northern fur seals: hypothesis testing supplants veri . cation . Journal of Parasitology 86, 1099-1106.
- Nguyen, KB & Smart, Jr , GC (1994). 14. Neosteinernema longicurvicauda n . gen., n. sp. (Rhabditia: Steinernematidae), a parasite of the termite Reticulitermes avipes (Koller). Journal of Nematology 26, 162-174.

VOLUME 04 ISSUE 01 Pages: 36-40

SJIF IMPACT FACTOR (2021: 5.478) (2022: 5.636) (2023: 6.741)

OCLC - 1368736135











- **15.** Nguyen, KB, Maruniak , J. & Adams, JB (2001). Diagnostic and phylogenetic utility of the rDNA internal transcribed spacer sequences of Steinernema. Journal of Nematology 33, 73-82.
- **16**. Poinar, Jr., GO (1986). Recognition of Neoaplectana species. Proceedings of the Helminthological Society of Washington 53, 121-129.
- Steiner, G. (1923). Aplectana Kraussei n. **17.** sp., eine in der Blattwespe Lyda sp. Nematodenform , nebst parasitic Bemerkungen über das Seitenorgan der parasitischen Nematoden. Zentralblatt fur Bacteriology, Parasitenkunde, Infections Krankheiten und Hygiene, Abteilung II 59, 14-18.
- 18. Steiner, G. (1929). Neoaplectana Glazer ngn sp. (Oxyuridae), a new nemesis parasite of the Japanese beetle (Popillia japonica Newm .). Journal of the Washington Academy of Science 19, 436-440.
- **19**. Stock, SP, Campbell, JF & Nadler, SA (2001).Phylogeny of Steinernema Travassos , 1927 (Cephalobina : Steinernematidae inferred) from ribosomal DNA sequences and morphological characters. Journal of Parasitology 87, 877-889.
- Stock, SP, Heng, J., Hunt, DJ, Reid, AP, Shen 20. , X. & Choo, HY (2001). Redescription of Steinernema longicaudum Shen & Wang (Steinernematidae Nematoda : geographic distribution and phenotypic

- variation between allopatric populations. Journal of Helminthology 75, 81-92.
- 21. 348. Stock, SP, A. Kusakabe, and RA Orozco, Secondary Metabolites Produced by. J Nematol, 2017. 49 (4): p. 373-383.
- 22. Travassos, L. (1927). Sobre o genero Oxysomatium . My ticket Biologico , Sao Paulo 5, 20-21.
- 23. Kurbonova NS (2023): Can entomopathogenic nematodes be the best for measure growing eco-friendly agricultural products? American Journal of Applied Science and Technology. ISSN-2771-2745. Sjifimpact Factor (2023: 7.063). Issue: Vol. 3 No. 07: Volume 03 Issue 07 | Pages: 23-31 Crossref DOI: https://doi.org/10.37547/ajast/Volume0 3 Issue07-06.
- 24. Kurbonova NS (2023): First report on local entomopathogenic nematode Steinernema feltiae in Uzbekistan. International Journal of Advance Scientific Research, 225-235. 3(07), https://doi.org/10.37547/ijasr-03-07-38